

Please hand in answers no later than **Friday 19th November**.

(Question 1) Consider the DS $x_{n+1} = F_\nu(x_n)$ where

$$F_\nu(x_n) = \nu + x^2,$$

where $x \in \mathbb{R}$ and $\nu \in \mathbb{R}$.

- (a) Find the fixed points. For what range of values of ν do they exist?
- (b) Find the value of ν for which there is a saddle-node bifurcation.
- (c) Find the value of ν for which there is a flip bifurcation. Is it super- or subcritical?
- (d) Sketch the bifurcation diagram in the (ν, x) plane; indicate the stability of the fixed points in your diagram.

(Question 2) Consider the DS $x_{n+1} = H_\mu(x_n)$ with

$$H_\mu(x) = \mu \tan^{-1} x,$$

where x is a real variable and μ is a real parameter.

- (a) How many fixed points are there? Specify the ranges of values of μ for which they exist.
- (b) Calculate the Schwarzian derivative of H_μ .
- (c) Describe the bifurcations which occur for
 - i. $\mu = 1$,
 - ii. $\mu = -1$.

If there are flip bifurcations, state whether they are supercritical or subcritical.

- (d) Sketch the bifurcation diagram in the (μ, x) plane. Indicate the stability of the fixed points in your diagram.